

SUPERSYMMETRIC DOMAIN WALLS (NEW EXACT RESULTS IN SUPERSYMMETRIC GLUODYNAMICS)

M. SHIFMAN

Theoretical Physics Institute, Univ. of Minnesota, Minneapolis, MN 55455, USA

Supersymmetric domain walls are discussed. The main emphasis is made on saturated walls in strongly coupled gauge theories. The central charge of the $\mathcal{N} = 1$ superalgebra appears in supersymmetric gluodynamics as a quantum anomaly. I also consider some rather unusual general features of supersymmetric domain walls.

The subject of my talk today lies somewhat outside the mainstream research in supersymmetry (SUSY). On one hand, it has little to do with the supersymmetric phenomenology, and, on the other hand, it does not blend with the current “high theory” – D -branes – although in a couple of points there is a certain overlap. Nevertheless, I think that the supersymmetric domain wall is a beautiful object that deserves thorough investigation. As we will see, this is a novel theoretical construction which admits exact results in strongly coupled theories, such as supersymmetric gluodynamics, the theory of gluons and gluinos, the closest relative of QCD. It is assumed that color confinement takes place in the strong coupling regime, much in the same way as in QCD. Usually, it is believed that under the circumstances no interesting dynamical quantity can be calculated exactly. I will explain that the domain wall tension is *exactly* calculable. Although my prime concern is SUSY gluodynamics, some of the results I am going to report will be of a more general nature. They refer to any SUSY theory with multiple domain walls.

When G. Dvali and I started working on the supersymmetric domain walls at the end of 1996^{1,2}, I could hardly anticipate that the project would grow and branch to the extent it actually did. Quite a few papers were published^{1–7}, and more are in progress. It goes without saying that it is impossible to present a full account of all the results obtained. I will limit myself to main findings; they are summarized below:

- It turns out that in $\mathcal{N} = 1$ theories, SUSY algebras may have central extensions. This assertion by itself is heresy. If you open any text book on supersymmetry you will read that two theorems forbid central extensions in $\mathcal{N} = 1$ superalgebras. I will explain how these theorems are circumvented in those cases in which domain walls are formed.
- A zoo of various domain walls is typical of SUSY theories possessing a discrete set of vacua. I keep in mind that supersymmetry guarantees the vanishing of the vacuum energy density, so multiple degenerate (and physically

inequivalent) vacua are quite typical.

- In certain domain walls, 1/2 of the supersymmetry is spontaneously broken and 1/2 preserved. These are the so called BPS walls (more exactly, they should be called BPS-saturated). The exact results mentioned at the beginning refer to the BPS walls. For such walls the energy density per unit area (the tension) is exactly calculable: it is expressed as the difference of the values of the superpotential far to the left and far to the right from the wall.

- In supersymmetric gluodynamics the role of the superpotential is played by the gluino condensate. This is not seen at the classical level. The central charge in SUSY gluodynamics vanishes at the classical level; it emerges at the quantum level as a quantum anomaly. The wall tension ε is

$$\varepsilon = \frac{N_c}{8\pi^2} |\Delta(\text{Tr } \lambda\lambda)|, \quad (1)$$

where N_c is the number of colors and λ is the gluino field. The internal structure of the domain wall may be incredibly complicated. Still, its tension is given by this very simple expression. I should add that the gluino condensate, in turn, was exactly calculated in 1988⁸. The fact that the gluino condensate is akin to the superpotential was known previously in a different context⁹. The correspondence gets an independent confirmation with the calculation of the central charge in SUSY gluodynamics. The central charge turns out to be proportional to $\lambda\lambda$.

- For certain domain walls there exists an exact degeneracy: say,

$$\varepsilon_{13} = \varepsilon_{12} + \varepsilon_{23} \quad (2)$$

where ε_{ij} is the tension of the wall interpolating between the vacuum i and j , respectively. In other words, if one puts the two domain walls indicated on the right-hand side at an arbitrary distance from each other, they absolutely do not interact, either classically or quantum mechanically, there is neither repulsion nor attraction. I believe that previously such domain walls were not known (although in some exactly solvable two-dimensional models a similar property was known for kinks).

After this brief “executive summary” I will move on to a more systematic discussion of the issues. First of all, how can one understand the existence of the central extensions of $\mathcal{N} = 1$ superalgebras? A heuristic explanation was suggested by Witten¹⁰. Assume a four-dimensional theory admits domain walls (or strings). Compactify the theory in two directions parallel to the wall on a very large two-torus of area A . The theory then reduces to an effective two-dimensional theory where the central extension obviously exists¹¹; the central charge for the states interpolating between distinct vacua at spatial

infinities is the superpotential difference between the two vacua. On the other hand, in the limit $A \rightarrow \infty$ we return back to the original four-dimensional theory.

Thus, there is no central charge for particles in four-dimensional $\mathcal{N} = 1$ theories, but either strings or domain walls can be BPS-saturated (i.e. have their mass equal to the central charge).

What allows one to circumvent the classic analysis of Haag *et al.*¹² asserting the impossibility of the central extension in $\mathcal{N} = 1$ theories? Implicit in this analysis is the assumption that the full four-dimensional Lorentz invariance is unbroken. Needless to say that in the sector with the given domain wall the Lorentz symmetry is spontaneously broken. The bosonic zero mode on the wall is the manifestation of this breaking. It is only the three-dimensional Lorentz invariance that survives (one time plus two spatial directions parallel to the wall).

The actual calculation of the central charge is quite trivial in the generalized Wess-Zumino models where it appears at the tree level^{1,4}. One calculates $\{Q_\alpha Q_\beta\}$ using the canonic commutation relations. Terms that are spatial integrals of total derivatives, that are usually dropped, must be kept, since the central charge emerges as a spatial integral of a total derivative. This is the only relatively subtle point. The result is not worth presenting here since shortly I will give a more general formula valid in gauge theories with arbitrary superpotentials.

The basis of any gauge theory is supersymmetric gluodynamics, and its Lagrangian is

$$\mathcal{L} = \left\{ \frac{1}{4g_0^2} \text{Tr} \int d^2\theta \ W^2 + \text{H.c.} \right\} = \frac{1}{g_0^2} \left\{ -\frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a + i \bar{\lambda}_\alpha^a \mathcal{D}^{\dot{\alpha}\alpha} \lambda_\alpha^a \right\}. \quad (3)$$

Note that in my normalization the coupling constant is an overall factor in front of $\text{Tr} W^2$. For simplicity I will limit myself to $\text{SU}(N)$ gauge groups, although the analysis can be readily made general.

Superficially, this theory is very similar to QCD – the only distinction is the representation of the fermion fields. In QCD the fermion fields belong to the fundamental representation, while in supersymmetric gluodynamics they are the Weyl fields in the adjoint representation. The theory is strongly coupled. It is believed that the theory confines color more or less in the same way as QCD. I hasten to add that, perhaps, this is not quite true, as will be explained shortly.

Supersymmetric gluodynamics has multiple discrete vacua. The Lagrangian (3) is invariant under $\text{U}(1)$ which is broken by the anomaly down to Z_{2N} . The expectation value $\langle \lambda^2 \rangle$ is the order parameter; the gluino condensation implies

that the discrete chiral symmetry Z_{2N} is spontaneously broken down to Z_2 ,

$$\langle \lambda^2 \rangle = \Lambda^3 \exp(2\pi i k/N), \quad k = 0, 1, 2, \dots, N-1. \quad (4)$$

The gluino condensation was conjectured by Witten in the early 1980's¹³. The value of the gluino condensate was exactly calculated⁸ by adding matter in an appropriate representation, Higgsing the theory – to enforce the weak coupling regime – and then returning back to SUSY gluodynamics by exploiting holomorphy.

The number of vacua labeled by the gluino condensate (4) is N . This is precisely the value of Witten's index for the $SU(N)$ SUSY gluodynamics. Thus, the chirally asymmetric vacua saturate Witten's index. Recently it was argued¹⁴ that the theory admits also a chirally symmetric vacuum ($\langle \lambda^2 \rangle = 0$) in which no mass gap develops. In this vacuum, the theory is presumably in the conformal regime; the energy density becomes positive in finite volume, so that the chirally symmetric vacuum does not contribute to Witten's index. It plays a role, however, in the domain wall dynamics.

Thus, we have multiple vacua which are degenerate. Supersymmetry guarantees that the vacuum energy density vanishes in each of these states. If so, there must exist field configurations interpolating between distinct vacua at infinities. By analogy with solid state physics, one calls them the domain walls. Assume the domain wall lies in the xy plane. At $z = -\infty$ the order parameter takes some value, say, $\lambda^2 = -\Lambda^3$. The value of λ^2 rapidly changes in the transition domain near $z = 0$ (the width of the transition domain is of order Λ^{-1}), then λ^2 approaches $+\Lambda^3$ at $z = +\infty$. Inside the transition domain the fields have to restructure themselves – there are gradient terms, and potential terms – so that the vacuum energy density is distorted and is of order Λ^4 . The total wall mass is proportional to $A\Lambda^3$, where A is the wall area, $A \rightarrow \infty$. The wall tension ε (i.e. the energy per unit area) is finite. Since the theory is strongly coupled we have no idea how to describe the internal structure of the wall. Yet, the tension is calculable. If you wish, the wall is a dynamic realization of 2-branes that are so popular now. The D -branes are abstract creatures, however, while the walls I am speaking about are made from “flesh and blood”.

In order to explain how one can exactly calculate ε , I must take a step back and discuss the superalgebra. The standard $\mathcal{N} = 1$ superalgebra includes two basic anticommutation relations,

$$\{Q_\alpha, \bar{Q}_{\dot{\beta}}\} = 2P_{\alpha\dot{\beta}} \quad \text{and} \quad \{\bar{Q}_{\dot{\alpha}}, \bar{Q}_{\dot{\beta}}\} = 0. \quad (5)$$

The vanishing of the anticommutator $\{\bar{Q}\bar{Q}\}$ is a consequence of the Lorentz symmetry and the Coleman-Mandula theorem¹⁵. At the classical level Eq.

(5) *is valid* in SUSY gluodynamics, as can be seen from a straightforward calculation of the canonic commutators. However, at the loop level a quantum anomaly modifies this relation. In a generic SUSY gauge theory with a superpotential $\mathcal{W}$ and a set of the matter fields in the representations R_i of the gauge group, one has^{2,3,4}

$$\{\bar{Q}_{\dot{\alpha}}\bar{Q}_{\dot{\beta}}\} = \frac{4}{3}(\vec{\sigma})_{\dot{\alpha}\dot{\beta}} \int d^3x \vec{\nabla} \left\{ \left[3\mathcal{W} - \sum_i Q_i \frac{\partial \mathcal{W}}{\partial Q_i} \right] - \left[\frac{3T(G) - \sum_i T(R_i)}{16\pi^2} \text{Tr } W^2 + \frac{1}{8} \sum_i \gamma_i \bar{D}^2 (\bar{Q}_i^+ e^V Q_i) \right] \right\}_{\theta=0} \quad (6)$$

Here $(\vec{\sigma})_{\dot{\alpha}\dot{\beta}}$ is a triplet of matrices transforming the vectorial index of the $\{0, 1\}$ representation of the Lorentz group into the spinorial indices, $(\vec{\sigma})_{\dot{\alpha}\dot{\beta}} = \{-\tau_3, i, \tau_1\}_{\dot{\alpha}\dot{\beta}}$. Moreover, $T(R)$ is (one half) of the Dynkin index for representation R , $T(G)$ is (one half) of the Dynkin index in the adjoint representation ($T(G) = N$ for $\text{SU}(N)$), γ_i are the anomalous dimensions of the matter fields Q_i .

The central extension in the right-hand side of Eq. (6) is proportional to the spatial integral of a total derivative. Thus, it vanishes for all localized field configurations (particles!), in full accord with the classic theorems. It does not vanish, however, for wall-like configurations. In supersymmetric gluodynamics the central charge is proportional to the difference in the expectation values of the gluino condensate in the vacua between which the given wall interpolates (see Eq. (1)). When matter fields are added, the superpotential also enters the game, in a well-defined manner. If one drops total superderivatives in Eq. (6) (they have no impact on the central charges) the right-hand side can be rewritten as

$$4(\vec{\sigma})_{\dot{\alpha}\dot{\beta}} \int d^3x \vec{\nabla} \left\{ \mathcal{W} - \frac{T(G) - \sum_i T(R_i)}{16\pi^2} \text{Tr } W^2 \right\}. \quad (7)$$

One can dub the combination in the braces a *generalized superpotential*. That's what replaces superpotential when the anomaly is switched on.

Do we deal with a new anomaly? Taking into account the geometric nature of the supercharges, the occurrence of a new anomaly seems highly unlikely. And, indeed, one can show⁴ that the right-hand side of Eq. (6) is in one-to-one correspondence with well-known anomaly¹⁶ in the superdivergence of the supercurrent¹⁷, $\bar{D}^{\dot{\alpha}} J_{\alpha\dot{\alpha}}$.

The next question to ask is, what are the consequences of central extension? In general, this question was investigated about 20 years ago¹¹. It was

shown that the nonvanishing central charge implies a lower bound on the mass of the soliton – the domain wall in the case at hand. If the lower bound is saturated (see e.g. Eq. (1) in SUSY gluodynamics), the wall is saturated; only 1/2 of supersymmetry is spontaneously broken on the wall, the other half is preserved. Thus, if the wall is saturated, we know exactly its tension, although not much is known about its internal structure.

Assume that the walls in SUSY gluodynamics are saturated (I will argue that this is the case later). Then the wall tension is given by Eq. (1). An intriguing question to study is the dependence on the number of colors. Apart from the explicit N_c factor on the right-hand side, we should keep in mind an implicit N_c dependence hidden in the gluino condensate. With my normalization of the fields, $\langle \text{Tr} \lambda^2 \rangle \sim N_c$; however, the variation of the gluino condensate in the neighboring vacua is much smaller. Indeed, from Eq. (4) one infers that for the neighboring vacua $\Delta \text{Tr} \lambda^2 \sim N_c^{-1} \text{Tr} \lambda^2 \sim N_c^0$. This means that the tension of such walls scales as $\varepsilon \sim N_c^1$, a rather unexpected dependence. The scaling law typical of glueball solitons is N_c^2 rather than N_c . Yet, the assumption of BPS saturation, in conjunction with the expression for the central charge we derived, unambiguously leads us to the linear N_c dependence. The very same linear dependence was observed by Witten who analyzed a theory related to SUSY gluodynamics from the point of view of D -brane physics¹⁰. In this picture the domain walls also appear naturally. The linear N_c dependence is natural from the D -brane perspective. The question is, can one understand the origin of the linear scaling law in field theory *per se*, without any reference to D -branes?

The answer to this question is “yes”⁷. At large N_c , the mesons and the glueballs should have masses of order 1, trilinear couplings of order $1/N_c$, and so on. This is conveniently encoded in an effective Lagrangian of the form

$$\mathcal{L} = N_c^2 F[M_i, \partial M_i], \quad (8)$$

where $\{M_i\}$ is a set of fields (generically infinite) representing mesons and glueballs. The value of the functional F itself and all its derivatives at the minima should be independent of N_c . This would ensure the proper N_c dependence of the masses and coupling constants. Now, suppose we have a solution of classical equations of motion M_{wall} which describes a wall configuration interpolating between two distinct minima. Since N_c^2 is an overall factor in Eq. (8), at first sight one may expect that $F[M_{\text{wall}}] = O(1)$, and, therefore, the wall tension $\varepsilon = O(N_c^2)$. Such a situation is standard in the soliton physics.

How can one avoid this conclusion? Consider a function F glued of N_c sectors, so that it is non-analytic. At the minima the derivative of F are N_c independent. If at distances of order $1/N_c$ in the space of fields M_i some of

the derivatives become large, of order N_c – which is typical of glued N_c -sector functionals – the naive N_c counting is distorted, and the overall N_c^2 factor in Eq. (8) is converted into N_c in the soliton mass.

Thus, the N_c -sector structure inherent to SUSY gluodynamics is an implicit source of N_c dependence. Of course, these sectors appear only in the “macroscopic” glueball-based description of the theory (8). In the “microscopic” description everything is smooth. The fact that the volume energy E is $O(N_c)$ inside the BPS wall connecting two neighboring vacua, say with $\text{Tr}\langle\lambda\lambda\rangle = \Lambda^3 \exp(2\pi i k/N)$ where $k = 0$ and 1 , is seen in the microscopic theory too. Indeed, for the BPS wall

$$G_{\mu\nu}^a G_{\mu\nu}^a \sim \partial_z \lambda^2 \sim \Delta \lambda^2 / L.$$

Since the volume energy density $E \sim N_c G_{\mu\nu}^a G_{\mu\nu}^a$, and $\Delta \lambda^2$ in the neighboring chirally asymmetric vacua is $O(1)$, we conclude that E scales as N_c .

Let me note in passing that the D -brane analysis¹⁰ suggests that the confining QCD string emanating from the probe color charges (quarks) on one side of the wall can terminate on the wall, without penetrating on the other side. In⁷ it is explained how this feature emerges in the field-theoretic picture of the domain walls.

Now, we pass to the most crucial point – whether or not the domain walls in SUSY gluodynamics are BPS saturated. This is a dynamical question. *A priori* it is impossible to assert that they are BPS saturated. A part of the walls – or all of them – could be nonsaturated. Then, Eq. (1) would play the role of a lower bound rather than the exact prediction.

We have very few tools which might help us to answer this question in SUSY gluodynamics. One can try to apply the Veneziano-Yankielowicz effective Lagrangian¹⁸, amended in¹⁴ to properly incorporate the non-anomalous Z_{N_c} symmetry of SUSY gluodynamics. Of course, since this Lagrangian is not truly Wilsonian, one cannot trust it in the quantitative aspect. However, since saturation *versus* non-saturation is a qualitative issue – the answer depends only on a global “geography” in the space of fields – it is reasonable to expect that the Veneziano-Yankielowicz Lagrangian can be applied for this limited purpose. Is this expectation true or false? Both...

The Lagrangian realizing all anomalous Ward identities is written in terms of the chiral superfield $S = 3/(32\pi^2 N_c) \text{Tr } W^2$,

$$\mathcal{L} = \frac{1}{4} \int d^4\theta N_c^2 (\bar{S}S)^{1/3} + \frac{1}{3} \int d^2\theta N_c S (\ln S^{N_c} + 2\pi i n) + \text{H.c.} \quad (9)$$

From this Lagrangian you see that the potential energy is built from N_c

distinct sectors,

$$U(\phi) = 4N_c^2(\phi^*\phi)^{2/3} \ln(\phi e^{-i2\pi n/N_c}) \ln(\phi^* e^{i2\pi n/N_c}) \quad (10)$$

$$\text{at } \frac{(2n-1)\pi}{N_c} < \arg\phi < \frac{(2n+1)\pi}{N_c}.$$

The Z_{N_c} symmetry is explicit in this expression. Including the kinetic term one gets the following effective Lagrangian:

$$\mathcal{L} = N_c^2 \{ \partial_\mu \phi^{1/3} \partial_\mu \phi^{*1/3} + U(\phi) \}. \quad (11)$$

It is quite obvious that we deal here with $N_c + 1$ supersymmetric minima – N_c minima at $\phi = e^{i2\pi n/N_c}$, corresponding to a non-vanishing value of the gluino condensate (spontaneously broken discrete chiral symmetry), and a minimum at $\phi = 0$ (unbroken chiral symmetry). The scalar potential itself is continuous, but its first derivative in the angular direction experiences a jump at $\arg\phi = (2n+1)\pi/N_c$.

Equations (10) and (11) do imply that the wall connecting any of the chirally asymmetric vacua with the chirally symmetric one at the origin is saturated. Building this wall is more or less a straightforward exercise³. At the same time, the walls connecting the chirally asymmetric vacua with each other are not seen, at least at $N_c = 3$ ¹⁹. In fact, they are not expected to be seen, since whenever a cusp is crossed, the description based on the Veneziano-Yankielowicz Lagrangian fails⁷. The restructuring of degrees of freedom responsible for the cusp is not properly reflected. Thus, in this case, the existence of the BPS walls remains to be confirmed. Witten's D brane construction¹⁰ can be viewed as an indirect argument and a guideline.

Since the chirally symmetric vacuum plays such an important role in this range of questions, it is worth discussing in more detail. Apart from the Veneziano-Yankielowicz Lagrangian there is one more argument (I call it the *missing state argument*) pointing in this direction. It is based on the comparison of two alternative calculations of $\langle \text{Tr}\lambda^2 \rangle$. One is the direct one-instanton calculation in the strong coupling regime. Another calculation uses the trick I mentioned at the beginning of my talk – the theory is expanded by adding matter fields which set up the Higgs regime. This is a weak coupling calculation, believed to be absolutely clean. The two results do not match²⁰! One can argue that the instanton calculation in the strong coupling regime averages the gluino condensate over all vacuum states²¹. If so, the existence of the chirally symmetric vacuum can be inferred through a leakage in this vacuum which would explain the mismatch.

The chirally symmetric vacuum at $\langle \text{Tr}\lambda^2 \rangle = 0$ is not seen in Witten's D -brane construction. Perhaps, this is not surprising at all. Indeed, there is

a good deal of extrapolation in this construction, against which the chirally symmetric vacua are stable (they have no choice since they have to saturate Witten's index) while the $\langle \text{Tr} \lambda^2 \rangle = 0$ vacuum need not be stable. In fact, as I mentioned, it disappears in finite volume. Evidently, it cannot survive the space-time distortions associated with the D -brane engineering.

Concluding my talk, I would like to make two brief remarks regarding the topics I had no time to cover. Both remarks do not necessarily refer to gauge theories; they are of a more general nature.

First, supersymmetric domain walls give a very interesting example of a degeneracy which has no analogs in the nonsupersymmetric case. Consider a theory with, say, three discrete vacua, such that the values of the superpotential in all vacua have one and the same phase. (In the gauge theories one should consider a generalized superpotential, see Eq. (7)). Such a situation is quite common in the Wess-Zumino models with two or more fields. Then $C_{13} = C_{12} + C_{23}$ where C 's denote the corresponding central charges. If BPS walls exist in all possible transitions, 12, 13 and 23, the wall tensions will obey a similar relation, see Eq. (2). This implies, in turn, that the 12 and 23 walls at rest do not interact with each other no matter how large/small the distance is between them^{5,6}. In a two-field Wess-Zumino model, it is possible to find an analytic solution describing a superposition of the 12 and 23 walls, for the first time ever⁶.

Second, one can use supersymmetric domain walls and similar topological defects in the cosmological aspect, as a possible mechanism for breaking SUSY, in part or completely, at scales much lower than M_{Planck} . It is difficult to overestimate the value of possible alternative scenarios of SUSY breaking occurring at a human scale. The topic was open for discussion in Ref. 1 and was further elaborated in²² where a relatively viable and even attractive scenario was put forward.

Acknowledgments

This work was supported in part by DOE under the grant number DE-FG02-94ER40823.

1. G. Dvali and M. Shifman, *Nucl. Phys.* **B504** (1997) 127.
2. G. Dvali and M. Shifman, *Phys. Lett.* **B396** (1997) 64; (E) **B407** (1997) 452.
3. A. Kovner, M. Shifman, and A. Smilga, *Phys. Rev.* **D56** (1997) 7978.
4. B. Chibisov and M. Shifman, *Phys. Rev.* **D56** (1997) 7990.
5. M. Shifman, *Phys. Rev.* **D57** (1998) 1258.

6. M. Shifman and M. Voloshin, *Phys. Rev.* **D57** (1998) 2590.
7. I. Kogan, A. Kovner, and M. Shifman, *Phys. Rev.* **D57** (1998) 5195.
8. M. Shifman and A. Vainshtein, *Nucl. Phys.* **B296** (1988) 445.
9. K. Intriligator and N. Seiberg, *Nucl. Phys. Proc. Suppl.* **45BC** (1996) 1 [hep-th/9509066].
10. E. Witten, *Nucl. Phys.* **B507** (1997) 658.
11. E. Witten and D. Olive, *Phys. Lett.* **B78** (1978) 97.
12. R. Haag, J. Lopuszanski and M. Sohnius, *Nucl. Phys.* **B88** (1975) 257.
13. E. Witten, *Nucl. Phys.* **B202** (1982) 253.
14. A. Kovner, and M. Shifman, *Phys. Rev.* **D56** (1997) 2396.
15. S. Coleman and J. Mandula, *Phys. Rev.* **159** (1967) 1251.
16. M. Grisaru, in *Recent Developments in Gravitation* (Cargèse Lectures, 1978), Eds. M. Levy and S. Deser (Plenum Press, New York, 1979), page 577, and references therein.
17. S. Ferrara and B. Zumino, *Nucl. Phys.* **B87** (1975) 207.
18. G. Veneziano and S. Yankielowicz, *Phys. Lett.* **113B** (1982) 231.
19. A. Smilga and A. Veselov, *Phys. Rev. Lett.* **79** (1997) 4529; *Nucl. Phys.* **B515** (1998) 163; hep-th/9801142; A. Smilga, hep-th/9711032.
20. V. Novikov, M. Shifman, A. Vainshtein, and V. Zakharov, *Nucl. Phys.* **B260** (1985) 157 (see App. B).
21. D. Amati *et al.*, *Phys. Reports*, **162** (1988) 169.
22. N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, *Phys. Lett.* **B429** (1998) 263; hep-ph/9807344.